



Substrate-driven microbial response: A novel mechanism contributes significantly to temperature sensitivity of N₂O emissions in upland arable soil



Alin Song^a, Yongchao Liang^b, Xibai Zeng^c, Huaqun Yin^d, Duanyang Xu^e, Boren Wang^a, Shilin Wen^a, Dongchu Li^a, Fenliang Fan^{a,*}

^a Key Laboratory of Plant Nutrition and Fertilizer, Ministry of Agriculture, Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Sciences, Beijing 100081, China

^b Ministry of Education Key Laboratory of Environment Remediation and Ecological Health, College of Environmental and Resource Sciences, Zhejiang University, Hangzhou 310058, China

^c Institute of Environment and Sustainable Development in Agriculture, Chinese Academy of Agricultural Sciences, Beijing 100081, China

^d Key Laboratory of Biometallurgy of Ministry of Education, School of Minerals Processing and Bioengineering, Central South University, Changsha 410083, China

^e Institute of Geographic Sciences and Natural Resources Research, CAS, Beijing 100101, China

ARTICLE INFO

Keywords:

Nitrous oxide
Temperature sensitivity
Microbial response
Nitrification
Denitrification
Anaerobic-zone development

ABSTRACT

The mechanism by which temperature sensitivity (TS) of soil N₂O emissions is increased by agricultural management with application of nitrogen fertilizer (AMN) is unclear. We hypothesized that a higher TS of N₂O emission induced by AMN is the result of the faster growth of specific microorganisms in response to faster nitrogen (N) mineralization at higher temperatures. To test this hypothesis, we used reciprocal transplants to separate the contributions of abiotic and microbial components to the TS of N₂O emissions in an arable soil receiving organic and inorganic fertilizers and its neighboring natural grassland soil treated with two levels of N. N₂O sources were separated with acetylene, and the abundances of N₂O-producing microbes were assessed by quantifying the copy numbers of the associated functional genes. Compared with natural soil, only changes in abiotic properties increased the Q₁₀ (the factor by which the rate increases with a 10 °C rise in temperature) by 105.7%, while changes in both abiotic and the microbiome increased the Q₁₀ by 225.2%. Higher TS of N₂O emission in the arable soil induced by a microbiome shift was associated with faster N mineralization, increased proportion of nitrification-N₂O emission, and faster growth of ammonia-oxidizing bacteria at higher temperatures. Addition of ammonium nitrate further enhanced the TS of N₂O emissions, the proportion of nitrification-N₂O emission, and the increased extent of the growth of ammonia-oxidizing bacteria in the soil with AMN compared to the natural grassland soil. Substrate-driven growth of ammonia-oxidizing bacteria with higher substrate preference contributes significantly to the higher TS of N₂O emission caused by AMN.

1. Introduction

Nitrous oxide (N₂O) is the third most important greenhouse gas (IPCC, 2007) and is a major cause of stratospheric ozone depletion (Ravishankara et al., 2009). About 60% of anthropogenic N₂O is emitted from agricultural soils, mostly derived from applications of synthetic fertilizer N and animal manure (Smith, 2017; Zhou et al., 2017). Besides by N sources, the emission of N₂O is controlled by many other factors, including contents of soil water and carbon and temperature (Weier et al., 1993; Conrad, 1996; Avrahami et al., 2003; Baggs and Philippot, 2010; Blagodatskaya et al., 2014; Smith, 2017). Of these, temperature is particularly significant because it influences

current N₂O emissions and also impacts future N₂O emissions via positive or negative feedback (Smith, 1997). We analyzed the results in the review by Smith (1997) and subsequent studies and found that N fertilization increases the N₂O-emission rate and also its temperature sensitivity (TS). The Q₁₀ levels in soils with AMN are higher than in soils without N fertilizer in both disperse studies and studies with paired N and no N treatments (Fig. 1, Table S1). However, the mechanisms underlying the AMN-enhanced TS of soil N₂O emission are not well known.

The variation in TS of soil N₂O emission has been attributed mostly to anaerobic-zone development (AZD) in aggregated soils (Smith, 1997, 2017). This explanation suggests that a temperature increase leads to an

* Corresponding author. South Zhongguancun Street No. 12, Beijing 100081, China.
E-mail address: fanfenliang@caas.cn (F. Fan).