Contents lists available at ScienceDirect

Field Crops Research

journal homepage: www.elsevier.com/locate/fcr

Quantification of yield gap and nutrient use efficiency of irrigated rice in China

Xinpeng Xu^{a,b}, Ping He^{a,c,*}, Shicheng Zhao^a, Shaojun Qiu^a, Adrian M. Johnston^d, Wei Zhou^{a,*}

^a Ministry of Agriculture Key Laboratory of Plant Nutrition and Fertilizer, Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Sciences (CAAS), Beijing 100081, PR China

^b Institute of Plant Nutrient and Resources, Beijing Academy of Agriculture and Forestry Sciences, Beijing 100097, PR China

^c International Plant Nutrition Institute (IPNI) China Program, CAAS-IPNI Joint Lab for Plant Nutrition Innovation Research, Beijing 100081, PR China

^d International Plant Nutrition Institute (IPNI), 102-411 Downey Road, Saskatoon, SK S7N4L8, Canada

ARTICLE INFO

Article history: Received 12 November 2015 Received in revised form 26 November 2015 Accepted 27 November 2015

Keywords: Rice Attainable yield Yield gap Yield response Nutrient use efficiency

ABSTRACT

Analyzing attainable yield (YA), yield gap (YG), and nutrient use efficiency (NUE) will help develop and inform agricultural policies and strategies to increase grain yield. Data from a total of 2218 on-farm rice experiments were collected between 2000 and 2013 from the main rice production areas of China. Common treatments in each study included the optimum nutrient management (OPT), farmers' fertilizer practices (FP) and nutrient omission treatments which were used to assess YG, yield response to nutrient (YR), and NUE. This study used meta-analysis and ANOVA to evaluate differences across the four rice planting seasons (early, middle, late, and single-season rice). The average YA from the OPT was 8.5 tha⁻¹, and the yield gap between OPT and FP was 0.6 tha⁻¹. The YR to nitrogen (N), phosphorus (P), and potassium (K) averaged 2.4, 0.9, and 1.0 tha⁻¹ across all sites, respectively. Results using the Pearson' correlation analysis showed significantly negative coefficients for YR to soil nutrient conditions and soil indigenous nutrient supplies. As compared to FP, the average recovery efficiency (RE) to N, P, and K with OPT increased by 10.1, 5.0 and 8.6 percent across all sites, respectively. In order to narrow the YG and increase NUE, effective soil, plant, nutrient management measures, advances in knowledge and technologies would be required to sustain higher crop production.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

Crop yield must increase substantially over the coming decades to keep pace with the increasing food demand of the world's population (Van Ittersum et al., 2013). Rice is an important staple food worldwide, and is the primary staple food in China. At present, the Chinese rice cultivation area is 30.3 million hectares with total rice

http://dx.doi.org/10.1016/j.fcr.2015.11.011 0378-4290/© 2015 Elsevier B.V. All rights reserved. yield reaching 203.6 million tons, and an average unit area yield of 6.7 tha^{-1} (China Agriculture Statistical Report, 2013). It is projected that world rice production needs to increase to 771.1 million tons by 2030 (Van Nguyen and Ferrero, 2006). Using aggregated data to quantify the yield gap (YG) on a regional, or agro-climatic zone, scale is helpful to quantify the scope for future yield improvement (Neumann et al., 2010; Van Wart et al., 2013).

Exploiting and narrowing the current yield gap (YG) between farmers practice and attainable yield by optimizing soil and nutrient management practices is one way to increase rice production. For sustainable intensification of rice cropping, it is important to identify areas with the greatest potential to narrow the YG, because there are large differences in climate, soil, and economic inputs among regions (Neumann et al., 2010; Van Ittersum et al., 2013). Many factors have contributed to the current YG, for example, the availability of production technologies, differences among rice varieties, a lack of balanced fertilizer application, and sub-optimal crop management by farmers. Many studies have shown that rice yield





CrossMark

Abbreviations: AE, agronomic efficiency; FP, farmers' practice; IKS, indigenous potassium supply; INS, indigenous nitrogen supply; IPS, indigenous phosphorus supply; NUE, nutrient use efficiency; OPT, optimum management; PFP, partial factor productivity; RE, recovery efficiency; YA, attainable yield; YG, yield gap; YR, yield response.

^{*} Corresponding authors at: Ministry of Agriculture Key Laboratory of Plant Nutrition and Fertilizer, Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Sciences (CAAS), Beijing 100081, PR China. Fax: +86 10 82106206.

E-mail addresses: phe@ipni.net, phe@caas.ac.cn (P. He), zhouwei02@caas.cn (W. Zhou).