

Vegetation patches increase wind-blown litter accumulation in a semi-arid steppe of northern China

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 Environ. Res. Lett. 11 124008

(<http://iopscience.iop.org/1748-9326/11/12/124008>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 80.82.78.170

This content was downloaded on 29/12/2016 at 04:26

Please note that [terms and conditions apply](#).

You may also be interested in:

[Modeling phenological responses of Inner Mongolia grassland species to regional climate change](#)

Qiuyue Li, Lin Xu, Xuebiao Pan et al.

[Grazing intensity and driving factors affect soil nitrous oxide fluxes during the growing seasons in the Hulunber meadow steppe of China](#)

Ruirui Yan, Huajun Tang, Xiaoping Xin et al.

[The physics of wind-blown sand and dust](#)

Jasper F Kok, Eric J R Parteli, Timothy I Michaels et al.

[Variation and control of soil organic carbon and other nutrients in permafrost regions on central Qinghai-Tibetan Plateau](#)

Wenjie Liu, Shengyun Chen, Qian Zhao et al.

[Elevated soil nitrogen pools after conversion of turfgrass to water-efficient residential landscapes](#)

Hannah Heavenrich and Sharon J Hall

[Drivers of tall shrub proliferation adjacent to the Dempster Highway, Northwest Territories, Canada](#)

Emily A Cameron and Trevor C Lantz

[Variations in soil carbon dioxide efflux across a thaw slump chronosequence in northwestern Alaska](#)

A E Jensen, K A Lohse, B T Crosby et al.

[Land cover/land use change in semi-arid Inner Mongolia: 1992–2004](#)

Ranjeet John, Jiquan Chen, Nan Lu et al.