



The effects of increased snow depth on plant and microbial biomass and community composition along a precipitation gradient in temperate steppes

Weixing Liu^a, Steven D. Allison^{b,c}, Ping Li^{a,d}, Jing Wang^{a,d}, Dima Chen^a, Zhenhua Wang^{a,d}, Sen Yang^{a,d}, Liwei Diao^a, Bin Wang^{a,d}, Lingli Liu^{a,d,*}

^a State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Xiangshan, Beijing 100093, China

^b Department of Ecology and Evolutionary Biology, University of California, Irvine, CA 92697, USA

^c Department of Earth System Science, University of California, Irvine, CA 92697, USA

^d University of Chinese Academy of Sciences, Yuquan Road, Beijing 100049, China

ARTICLE INFO

Keywords:

Microbial biomass
Microbial community
Plant biomass
Plant community
Plant-microbe relationship
Sensitivity

ABSTRACT

Shift in precipitation regime could greatly alter plant and microbial activity, and thus the contemporary and future ecosystem dynamics in grasslands. We investigated how changes in snow depth affect plants, microbes and their relationships after 10 consecutive years of snow treatments in different steppes. We selected 8 snow fences along a mean annual precipitation (MAP) gradient from 225 to 375 mm in Inner Mongolia. For each snow fence, study plots were set up at 7 transects with different levels of snow depth. We found that ecosystem properties, including soil moisture, the biomass and nitrogen (N) pools of microbes and plants, the fungi: bacteria ratio and the grass: forb ratio, increased with increasing snow depth at the drier sites with lower MAP, but not at the wetter sites with higher MAP. At any given site, the sensitivity of these ecosystem properties to changes in snow depth was determined by the slopes of these variables against snow depth. The results showed that the sensitivity of these ecosystem properties to changes in snow depth decreased linearly with increase in MAP levels. In addition, we also found that increased snow depth shifted the relationship between microbial and plant biomass from positive to negative. Our work reveals the importance of snow water in regulating plant and microbial processes in temperate steppes, especially under lower MAP conditions. The greater plant and microbial biomass and the shift of community toward greater fungi: bacteria and grass: forb ratio imply that increased snowmelt input alleviated water limitation in temperate steppes and altered plant and microbial communities. Our study helps to better predict that how changes in winter precipitation could affect the biomass and composition of plants and soil microbes in grasslands.

1. Introduction

Numerous studies have assessed how changes in precipitation regimes influence ecosystem structure and functioning using manipulation experiments (Bachar et al., 2010; Estiarte et al., 2016). Most of these studies focused on the effects of annual or summer precipitation on plant and microbial processes, and often with site-level data (Vicente-Serrano et al., 2013; Estiarte et al., 2016). Climate change is expected to alter the amount of precipitation falling as snow in Northern Hemisphere (Peng et al., 2010; IPCC, 2014). In addition, the changes in wind patterns or vegetation cover will also alter snow redistribution by wind drifting, especially in ecosystems with short-statured plants such as grasslands (Ayles et al., 2010). Snow could greatly affect water and nutrient cycles, especially during winter and early growing season (Schimel et al., 2004). However, we know very little about how

changes in snow depth or redistribution will affect the productivity and communities of plants and soil microbes, as well as their relationship across the precipitation gradient.

Changes in snow depth can affect soil microbial activity and plant growth through abiotic or biotic mechanisms (Buckeridge et al., 2010; Sorensen et al., 2016). Greater snow accumulation elevates soil temperature by insulating soils from cold winter air (Brooks et al., 2005; Natali et al., 2011), and increases soil moisture (Groffman et al., 2001). Aside from increasing water availability (Groffman et al., 2001), greater snow depth stimulates nitrogen (N) mineralization (Schimel et al., 2004; Freppaz et al., 2007, 2012) and increases N availability during spring, especially in N limited ecosystems (Buckeridge et al., 2010; Leffler and Welker, 2013). Together these mechanisms can increase water and nutrient supply for soil microbes and plants (Groffman et al., 2006; Buckeridge et al., 2010). In addition, greater snow accumulation

* Corresponding author. State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Xiangshan, Beijing 100093, China.
E-mail address: lingli.liu@ibcas.ac.cn (L. Liu).